Saskatchewan
Ministry of
Agriculture

Crop Report

Crop Report for the Period August 10 to 16, 2010

Published by the Ministry of Agriculture
ISSN 0701 7085
Report number 20, August 19, 2010

Just over two per cent of the 2010 crop is swathed or ready to straight-combine, according to Saskatchewan Agriculture's weekly Crop Report.

Less than one per cent of the crop has been combined. When broken down by crop, six per cent of the winter wheat and fall rye and three per cent of the lentils and peas have been combined. The five-year provincial average (2005 to 2009) for this time of year is five per cent combined. The wet and cool weather has delayed crop maturity and slowed harvest.

Haying continues. Rain, high humidity and heavy dew continue to slow progress. Saskatchewan livestock producers have 90 per cent of the hay crop cut, with 80 per cent of that baled or put into silage. Quality is rated as nine per cent excellent, 65 per cent good, 15 per cent fair and two per cent poor.

The estimated hay yield for both dryland alfalfa and alfalfa/brome stands is 2.1 tons per acre. Estimated yield on irrigated alfalfa and alfalfa/brome stands is 2.7 tons per acre. Greenfeed is expected to yield 2.4 tons per acre on dryland and three tons per acre on irrigated stands. The five-year average (2005 to 2009) yield on dry land alfalfa/brome stands is 1.4 tons per acre.

Cropland topsoil moisture is rated as 27 per cent surplus, 70 per cent adequate and three per cent short. Topsoil moisture on hay land and pasture is rated as 15 per cent surplus, 80 per cent adequate and five per cent short.

One year ago

Just over three per cent of the 2009 crop had been combined. Eleven per cent of the lentils and 10 per cent of the peas were swathed or were ready to straight combine. Haying was wrapping up.

Harvest progress was slow due to rain and cool weather.

The majority of crop damage is being caused by flooding, disease and hail.

Farmers are busy haying, getting ready for harvest and hauling grain.

Saskatchewan Crop Insurance Corporation Report

The deadline to select winterkill coverage for fall rye and winter wheat is August 25.

For further information, contact: Grant McLean, PAg,
Cropping Management Specialist, Agriculture Knowledge Centre,
Toll Free: 1-866-457-2377, E-mail: cropreport@gov.sk.ca.
Also available on the Ministry of Agriculture website at www.agriculture.gov.sk.ca.

Agriculture and Agri-Food Canada | Agriculture et Agroalimentaire Canada

SCIC
SASKATCHEWAN CROP INSURANCE CORPORATION

Southeastern Saskatchewan (Crop Districts 1, 2, and 3ASE)

All areas received between six and 60 mm of rain during the week. The Redvers area received 60 mm, the Gainsborough area 35 mm, the Maryfield area 45 mm, the Broadview area 9 mm, the Weyburn area 34 mm, the Wilcox area 50 mm, the Marquis area 32 mm, the Radville area 28 mm and the Ceylon area 14 mm.

The region has two per cent of the crop combined and over six per cent is swathed or ready to straight-cut. Twenty per cent of the winter wheat and fall rye, two per cent of the barley, 10 per cent of the lentils, nine per cent of the field peas and one per cent of the mustard have been combined. Twenty-nine per cent of the winter wheat and fall rye, four per cent of the spring wheat, seven per cent of the durum, one per cent of the oats, three per cent of the barley, two per cent of the lentils and 17 per cent of the field peas are swathed or ready to straight-combine.

Haying is proving difficult, with rain and high humidity slowing operations. Hay quality has also been reduced in some areas due to rain on the swath. Baling hours are cut short due humidity in the evening. Ninety-three per cent of the hay in the region is cut and 87 per cent of this has been baled or put into silage. Six per cent is in the swath. Quality is rated as seven per cent excellent, 73 per cent good and 13 per cent fair. Some producers are concerned that they may not get some hay fields cut.

Estimated hay yields are as follows: dryland alfalfa and alfalfa/brome stands, 2.3 tons per acre; dryland greenfeed, 1.9 tons per acre.

Some cattle producers are having issues with foot rot in cattle due to the wet conditions.

Cropland topsoil moisture conditions are reported as 18 per cent surplus, 78 per cent adequate, three per cent short and one per cent very short. On hay land and pasture, topsoil moisture is rated as 10 per cent surplus, 83 per cent adequate, five per cent short and two per cent very short. Twenty-two per cent of cropland in Crop District 3ASW is short of topsoil moisture.

Crop damage is mostly attributed to hail, wind and flooding. The following communities reported hail damage: Frobisher, Lampman, Manor, Tantallon, Fillmore and Ceylon. On August 12, wind and hail northwest of Manor damaged between 10 to 80 per cent of crops in a one mile by nine mile strip. There was also hail damage reported south of Carlyle. Disease is also causing crop damage. The recent rain is causing some disease concerns in ripening pulse crops. Sprouting and bleaching in swathed cereals is also a concern.

Crop maturity has slowed due to cool and wet weather. Some warm weather is needed to push the crops along. Many areas are still one to two weeks behind

normal in crop development. Farmers are haying, getting ready for harvest and controlling weeds on unseeded acres.

Southwestern Saskatchewan (Crop Districts 3ASW, 3AN, 3B and 4)

All areas of the region received between 12 and 82 mm of rain for the week. Most of the region reported above 20 mm the week. The Consul area received 82 mm, the Rockglen, Tyner and Gull Lake areas 65 mm, the Coderre area 39 mm, the Shaunavon area 42 mm and the Glenbain area received 12 mm.

The region has one per cent of the 2010 crop combined. Three per cent of the field peas, one per cent of the lentils, four per cent of the winter wheat and seven per cent of the fall rye have been combined. Sixteen per cent of the winter wheat, 31 per cent of the fall rye, one per cent of the oats, 12 per cent of the triticale, 10 per cent of the canola, two per cent of the mustard, four per cent of the lentils, 14 per cent of the field peas and three per cent of the chickpeas are swathed or are ready to straight-combine.

Haying is wrapping up in the region. Ninety-five per cent has been cut, and 89 per cent of that has been baled or put into silage. Quality is rated as 23 per cent excellent, 65 per cent good and four per cent fair. The estimated hay yield on dryland alfalfa and alfalfa/brome stands is two tons per acre. The estimated yield on irrigated alfalfa and alfalfa/brome stands is 2.8 tons per acre. Dryland greenfeed yield is estimated as 3.3 tons per acre.

Cropland topsoil moisture is rated as 13 per cent surplus, 81 per cent adequate and six short. Hay land and pasture topsoil moisture conditions are rated as five per cent surplus, 86 per cent adequate and nine per cent short. CDs 3BN and 4B are reporting 29 and 22 per cent surplus topsoil moisture, while CD 3ASW is reporting 22 per cent short topsoil moisture on cropland.

Crop damage has been caused by hail, disease and flooding. Hail damage was reported in the following areas: Rockglen, Glentworth, Consul, Shaunavon, Gull Lake and Maple Creek. Disease is also causing damage to pulses.

Farmers are busy finishing haying, getting ready for harvest, hauling bales and scouting fields. The recent rain has farmers with desiccated pulses and swathed cereals worried about crop quality. Farmers are also concerned about an early frost, as nighttime temperatures are dropping and crops are two weeks behind in most cases. Crops are lodged in areas where there were heavy rains. A few weeks of dry and sunny weather are needed.

East-Central Saskatchewan (Crop Districts 5 and 6a)

All of the region's crop reporters recorded rain in amounts ranging from four to 80 mm. Since April 1, between 259 and 716 mm of rain has fallen on the region. The Saltcoats area received 75 mm, the Ituna area 4 mm, the Elfros area 80 mm, the Foam Lake area 70 mm, the Quill Lake area 22 mm, the Lumsden area 9 mm, the Allan area 64 mm and the Craik area 18 mm.

One per cent of the fall rye has been combined. Twelve per cent of the winter wheat, 31 per cent of the fall rye, one per cent of the barley and three per cent of the field peas are swathed or ready to straight-combine.

Haying is continuing, albeit slowly due to high humidity and rain showers. Eighty-eight per cent of the hay is cut, and 75 per cent of that has been baled up or put into silage. Quality is rated as eight per cent excellent, 60 per cent good and 20 per cent fair. The yield on dryland alfalfa and alfalfa/brome stands is estimated to be 2.2 tons per acre. Dry land green feed yield is estimated to be 2.1 tons per acre.

Topsoil moisture on cropland is reported as 44 per cent surplus, 54 per cent adequate and two per cent short. Hay land and pasture topsoil moisture is rated as 29 per cent surplus, 68 per cent adequate and three per cent short. CD 5B is reporting 63 and 49 per cent surplus topsoil moisture on cropland and hay land and pasture, respectively.

Flooding, disease and hail were the major causes of crop damage. The Churchbridge area reported flooding. Hail damage was reported in the Churchbridge, Stockholm and Saltcoats areas. Disease in pulse crops and lodging in areas that received heavy rain are causing concerns as well.

Farmers are busy haying and getting ready for harvest. Early-seeded wheat and pulses are being desiccated and swathed. Crops need two to three weeks to mature. Sunny and dry weather is needed. The land has not had any opportunity to dry out. Getting into the wet fields with equipment is very difficult.

West-Central Saskatchewan (Crop Districts 6B and 7)

All areas reported between six to 79 mm of rain for the week. Most areas received more than 25 mm of moisture. The Conquest, Netherhill and Dinsmore areas received 66 mm, the Sonningdale area 6 mm, the Herschel area 79 mm, the Kerrobert area 49 mm and the Cando and Battleford area 13 mm.

Three per cent of the winter wheat and one per cent of the fall rye have been combined. Thirty-six per cent of the winter wheat, 31 per cent of the fall rye and one per cent of the canola and mustard is swathed or ready to straight-combine.

Haying continues. The hay swaths are slow to dry due to rain and high humidity. Most of the hay that has been cut in the past week has been rained on at least

once. Eighty-three per cent of the hay crop has been cut, and 70 per cent of that has been baled or put into silage. Quality is rated as 63 per cent good, 26 per cent fair and five per cent poor. The estimated yield on dryland alfalfa and alfalfa/brome stands is 2.2 tons per acre. Dry land greenfeed is estimated to yield 2.6 tons per acre. The estimated yields on irrigated alfalfa and alfalfa/brome stands is 3.4 and 3.9 tons per acre, respectively.

Cropland topsoil moisture is rated as 33 per cent surplus and 67 per cent adequate. Hay land and pasture topsoil moisture conditions are rated as 24 per cent surplus and 76 per cent adequate.

The majority of crop damage is due to disease, flooding and hail. Due to recent moisture, farmers are concerned about quality as the crop matures. Pulses are showing high disease pressures, particularly in lentils, which is causing quality issues. Hail damage was reported in the Herschel, Smiley and Major areas.

Farmers are busy haying and getting ready for harvest. Crops are one to two weeks behind normal in development, so many farmers are hoping for warm, sunny weather and a long, open fall.

Northeastern Saskatchewan (Crop Districts 8 and 9AE)

All areas reported between 11 and 58 mm of rain for the week. The Porcupine Plain area received 46 mm, the Codette area 14 mm, the Alvena area 58 mm, the Humboldt area 38 mm and the Garrick area 23 mm.

Eight per cent of the winter wheat, 12 per cent of the fall rye, and one per cent of the field peas are swathed or ready to straight-combine. Canola is being swathed. Barley crops are ready to straight-cut in some areas.

Haying was interrupted again this week by rain, high humidity and heavy dew. Eighty-three per cent of the hay crop has been cut and 74 per cent of that has been baled or put into silage. Quality is rated as 12 per cent excellent, 75 per cent good and 13 per cent fair. Dryland hay yields are estimated to be 1.7 and 1.5 tons per acre on alfalfa and alfalfa/brome stands, respectively. Dryland green feed yield is estimated to be two tons per acre.

Cropland topsoil moisture is rated as 41 per cent surplus, 55 per cent adequate and four per cent short. Hay land and pasture topsoil moisture is rated as 42 per cent surplus, 52 per cent adequate and six per cent short. CD 8A is reporting 81 per cent surplus on crop land topsoil moisture.

Crop damage is due to flooding and disease. The heavy rain in some areas has lodged crops.

Farmers are busy haying and getting ready for harvest. Crop staging is variable across the region and crops are still one to two weeks behind normal in development. There are fields that are blooming. Many fields are still unable to support equipment. Most crop reporters are indicating it will be still two to three weeks until harvest is in full swing.

Northwestern Saskatchewan (Crop Districts 9AW and 9B)

All areas reported moisture in amounts ranging from nine to 68 mm. The Duck Lake area received 58 mm, the Spiritwood area 13 mm, the Neilburg and Meadow Lake areas 20 mm, the Lloydminster area 62 mm and the Rapid View area 4 mm.

Harvest has yet to begin in the region. Crops are starting to turn and peas are being desiccated.

Haying continues, but progress is slow due to rain, slow drying time and heavy dew. Some swaths have been in the field for two weeks. Quality of standing hay is starting to decline as it matures. Eighty-eight per cent of the hay crop is cut, with 69 per cent of that baled or put into silage. Quality is rated as 57 per cent good, 14 per cent fair and seven per cent poor. Hay yield on alfalfa and alfalfa/brome stands is estimated at 1.8 tons per acre. Dryland greenfeed yield is estimated at 2.1 tons per acre.

Cropland topsoil moisture is rated as six per cent surplus and 94 per cent adequate. Hay land and pasture topsoil moisture is rated as 100 per cent adequate.

Flooding, disease and hail were the causes of most of the crop damage. Hail damage was reported in the Radisson, Debden and North Battleford areas. Disease pressures in pulses have caused some crop damage as well. Some crops are lodging in areas that received heavy rain.

Two to three weeks of frost-free and sunny weather is needed to get the majority of the crop harvested. Farmers are busy haying, scouting fields and getting ready for harvest.

Crop Districts and Rural Municipalities in Saskatchewan



Weekly Rainfall Summary

(in millimeters) 1 inch = 25 mm for the period to August 10 to 16, 2010

Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr
1A	1	Argyle	35.0	417.0
	2	Mount Pleasant	N/A	285
	3	Enniskillen	36	329
	33 A	Moose Creek	N/A	249
	33 B	Moose Creek	N/A	323
	34	Browning	22	299
	61	Antler	60	490
	63	Moose Mountain	59	382
	64	Brock	21	349
	65	Tecumseh	12	361
1B	91	Maryfield	45	381
	92	Walpole	N/A	299
	93	Wawken	20	397
	96	Golden West	8	322
	122	Martin	N/A	390
	123 A	Silverwood	6	281
	123 B	Silverwood	N/A	N/A
	124	Kingsley	19	295
	125	Chester	8.9	300
	151 A	Rocanville	41	409
	151 B	Rocanville	N/A	409
	153	Willowdale	N/A	407
	154	Elcapo	9	395
	155	Wolseley	15	483
2A	66	Griffin	N/A	395
	67	Weyburn	34	394
	68	Brokenshell	25	318
	69	Norton	N/A	313
	96	Fillmore	8	288
	97	Wellington	17	243
2B	127 A	Francis	28	349
	127 B	Francis	11	232
	129	Bratt's Lake	50.5	284
	156 A	Indian Head	14	408
	156 B	Indian Head	21	388
	157	South Qu'appelle	N/A	NIL
	160	Pense	28	332
	161	Moose Jaw	25	340
	191	Marquis	32	394
3ASE	38 A	Laurier	28	373
	38 B	Laurier	15	311
	39 A	The Gap	14	277
	39 B	The Gap	20	427
3ASW	10	Happy Valley	14	354
	11	Hart Butte	N/A	NIL
	12	Poplar Valley	15	254
	40	Bengough	18	338
	42	Willow Bunch	10	321
	43	Old Post	65	317
	44	Waverley	38	352
	70	Key West	N/A	315
	71 A	Excel	N/A	293
	71 B	Excel	N/A	227
	73 A	Stonehenge	19.2	305
	73 B	Stonehenge	27	389
	74	Wood River	16	314
3AN	101	Terrell	24	371
	102	Lake Johnston	15.2	289
	103	Sutton	N/A	292
	134	Shamrock	39	397
	164	Chaplin	33.5	341
	193	Eyebrow	38	363
	194	Enfield	18	384
3BS	17	Val Marie	22	347
	45	Mankota	29	269
	75 A	Pinto Creek	N/A	289
	75 B	Pinto Creek	17	291
	76 A	Auvergne	22	267
	76 B	Auvergne	18	252
	77	Wise Creek	27	267
	78 A	Grassy Creek	N/A	331
	78 B	Grassy Creek	24	347
	78 C	Grassy Creek	N/A	332
	105	Glenbain	12	278
	106	Whiska Creek	37	234
	107	Lac Pelletier	N/A	274
	108	Bone Creek	42	361
3BN	138	Webb	53	371
	165	Morse	N/A	NIL
	166	Excelsior	29	466
	167	Sask. Landing	53.4	421
	168	Riverside	63	431
	226	Victory	N/A	372
	228	Lacadena	65	431
	256	Coteau	N/A	NIL
	257	Monet	33	387

Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr
4A	49	White Valley	33	333
	49 A	White Valley	31	269
	51	Reno	82	358
	79	Arlington	36	382
	109	Carmichael	N/A	466
	110	Piapot	N/A	353
	111	Maple Creek	N/A	355
4B	139	Gull Lake	65	486
	141	Big Stick	33	428
	142	Enterprise	73	514
	169	Pittville	43	380
	171	Fox Valley	41	379
	229	Miry Creek	63	494
	231	Happyland	13	406
5A	181	Langenberg	10	339
	183	Fertile Belt	32	393
	185	Mcleod	21	356
	186	Abernethy	16	302
	211	Churchbridge	16	368
	213	Saltcoats	75	410
	216	Tullymet	11	230
	241	Calder	16	336
	243	Wallace	N/A	335
	245 A	Garry	13	428
	245 B	Garry	15	473
	246	Ituna Bon Accord	4	305
	247	Kellross	18	433
	248	Touchwood	N/A	245
5B	271	Cote	26	351
	275	Insinger	N/A	NIL
	276	Foam Lake	N/A	562
	277	Emerald	70	585
	278	Kutawa	N/A	NIL
	305	Invermay	43	716
	307	Elfros	80	704
	308	Big Quill	34	451
	333	Clayton	N/A	134
	335	Hazel Dell	50	438
	336	Saaman	62	573
	337	Lakeview	43	453
	338	Lakeside	22	497
	366	Kelvington	41	394
6A	190 A	Dufferin	13	259
	190 B	Dufferin	17	358
	190 C	Dufferin	9	304
	190 D	Dufferin	25	339
	219 A	Longlaketon	42	314
	219 B	Longlaketon	43	360
	220	Mckillop	13	325
	221	Samia	25.4	411
	222	Craik	27	439
	251	Big Arm	44	419
	252	Arm River	18	353
	279	Mount Hope	70	424
	280	Wreford	N/A	323
	282	McCraney	31.5	432
	309	Prairie Rose	19	450
	313	Lost River	64	391
	339	Leroy	35.6	510
	340	Wolverine	44	507
	341	Viscount	49	481
	343	Blucher	55	500
6B	253	Willner	20	331
	283	Rosedale	61	503
	284	Rudy	36	477
	285	Fertile Valley	66	439
	286	Milden	66	457
	314	Dundurn	63	507
	344	Corman Park	N/A	430
	346	Perdue	56	522
	376 A	Eagle Creek	31	624
	376 B	Eagle Creek	6	452
7A	260	Newcombe	N/A	375
	260 B	Newcombe	13	293
	287	St. Andrews	34	364
	288	Pleasant Valley	24	379
	290 A	Kindersley	66	385
	290 B	Kindersley	N/A	349
	292	Milton	28	314
	317 A	Marriott	64	522
	317 B	Marriott	72	541
	318 A	Mountain View	79	496
	318 B	Mountain View	21	508
	320	Oakdale	53	351
	321	Prairiedale	24	376

Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr
7B	347 A	Biggar	16	405
	347 B	Biggar	N/A	415
	350 A	Mariposa	34	305
	350 B	Mariposa	49	373
	351	Progress	37	408
	352	Heart's Hill	26	437
	377	Glenside	24	458
	378 A	Rosemount	N/A	342
	378 B	Rosemount	13	434
	379	Reford	18	395
	380	Tramping Lake	18	350
	381	Grass Lake	73	494
	382	Eye Hill	31	386
	409	Buffalo	13	519
	410	Round Valley	23	437
	411	Senlac	30	416
8A	395	Porcupine	46	407
	397	Barrier Valley	44.4	478
	428	Star City	N/A	403
	456	Arborfield	35	383
	457	Connaught	N/A	521
	458	Willow Creek	14	449
	486	Moose Range	35	373
8B	369	St. Peter	42	478
	370	Humboldt	38	568
	371	Bayne	39	508
	372	Grant	36.6	480
	402	Fish Creek	58	416
	429 A	Flett's Springs	21	313
	429 B	Flett's Springs	46	456
	430	Invergordon	N/A	43
	459	Kinistino	29	417
9AE	461	Prince Albert	11	430
	488	Torch River	23	441
	520	Paddockwood	N/A	67
9AW	406	Mayfield	15	333
	435	Redberry	34	452
	436	Douglas	N/A	247
	463	Duck Lake	58	400
	464	Leask	22	448
	467	Round Hill	N/A	NIL
	494	Canwood	24	440
	496	Spiritwood	13	504
	497	Medstead	37	369
9B	438	Battle River	9	402
	440	Hillsdale	21.5	358
	442	Manitou Lake	20.3	330
	498 A	Parkdale	31	366
	498 B	Parkdale	N/A	296
	499	Mervin	36.6	314
	501 A	Frenchman Butte	38	343
	501 B	Frenchman Butte	19	306
	501 C	Frenchman Butte	66	419
	502	Britannia	68	431
	561	Loon Lake	32	337
	588 A	Meadow Lake	20	355
	588 B	Meadow Lake	4	370
	588 C	Meadow Lake	45	389
	588 D	Meadow Lake	19.8	365
	622	Beaver River	N/A	329

Municipality No: A, B, C, and D - more than one reporter

These precipitation amounts represent point locations within each municipality and do not necessarily reflect the whole R. M.

Weekly Rainfall

for the week ending August 16, 2010



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

© 2010 Government of Saskatchewan

0 25 50 100 150 200

Kilometers

N W E S

Projection: UTM Zone 13 Datum: NAD83

Data Source:
Rainfall - Ministry of Agriculture, Crop Report Database
Spline interpolation (tension = 50)

Prepared by: Geomatics Services Date: August 18, 2010

Weekly Rainfall

for the week ending August 16, 2010



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas this map should be used for regional analysis only

Saskatchewan Ministry of Agriculture

0 25 50 100 150 200

Kilometers

N

Data Source
Rainfall - Ministry of Agriculture, Crop Report Database
Spline interpolation (tension = 50)

Prepared by: Geomatics Services Date: August 18, 2

©2010 Government of Saskatchewan

Cumulative Rainfall

From: April 1, 2010
To: August 16, 2010



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

© 2010 Government of Saskatchewan

0 25 50 100 150 200

Kilometers

N W E S

Projection: UTM Zone 13 Datum: NAD83

Data Source:
Rainfall - Ministry of Agriculture, Crop Report Database
Spline interpolation (tension = 50)

Prepared by: Geomatics Services Date: August 18, 2010

Cropland Topsoil Moisture Conditions

August 17, 2010



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

© 2010 Government of Saskatchewan

0 25 50 100 150 200

Kilometers

N W E S

Projection: UTM Zone 13 Datum: NAD83

Data Source:
Moisture - Ministry of Agriculture, Crop Report Database
Spline interpolation (tension = 50)

Prepared by: Geomatics Services Date: August 18, 2010

Cumulative Rainfall

From: April 1, 2010
To: August 16, 2010



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

© 2010 Government of Saskatchewan

0 25 50 100 150 200

Kilometers

N W E S

Projection: UTM Zone 13 Datum: NAD83

Data Source:
Rainfall - Ministry of Agriculture, Crop Report Database
Spline interpolation (tension = 50)

Prepared by: Geomatics Services Date: August 18, 2010

Cropland Topsoil Moisture Conditions

August 17, 2010



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

©2010 Government of Saskatchewan

0 25 50 100 150 200

Kilometers

N W E S

Projection: UTM Zone 13 Datum: NAD83

Data Source:
Moisture - Ministry of Agriculture, Crop Report Database
Spline interpolation (tension = 50)

Prepared by: Geomatics Services Date: August 18, 2010

Hay and Pasture Topsoil Moisture Conditions

August 17, 2010



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

© 2010 Government of Saskatchewan

0 25 50 100 150 200

Kilometers

N W E S

Projection: UTM Zone 13 Datum: NAD83

Data Source:
Moisture - Ministry of Agriculture, Crop Report Database
Spline interpolation (tension = 50)

Prepared by: Geomatics Services Date: August 18, 2010

Hay and Pasture Topsoil Moisture Conditions

August 17, 2010



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Saskatchewan Ministry of Agriculture

© 2010 Government of Saskatchewan

0 25 50 100 150 200

Kilometers

N W E S

Projection: UTM Zone 13 Datum: NAD83

Data Source:
Moisture - Ministry of Agriculture, Crop Report Database
Spline interpolation (tension = 50)

Prepared by: Geomatics Services Date: August 18, 2010